



Please be advised that this resource may contain images of Aboriginal or Torres Strait Islander people who are now deceased.

Acknowledgements

This resource has been developed in consultation with Aboriginal people, health specialists, health organisations and community groups. We are thankful to all who contributed.

The development of this booklet has been informed by the 'Having a Healthy Baby' booklets for women. The 'Having a Healthy Baby' booklets were produced as part of the National Diabetes Services Scheme (NDSS) Diabetes in Pregnancy National Development Program. The NDSS is an initiative of the Australian Government administered by Diabetes Australia. Copyright in the 'Having a Healthy Baby' booklets belongs to the Australian Government.

Diabetes WA would like to thank the Aboriginal women and their families, and the Aboriginal Health Practitioners/Workers who kindly donated their time to be photographed for this booklet.

This booklet will help you understand gestational diabetes. The information in this booklet should not take the place of advice you will receive from your doctor and diabetes specialists. It aims to help you to learn as much as you can about gestational diabetes so that you can have a healthy pregnancy and a healthy baby.


Contents

What is gestational diabetes?	4
What causes gestational diabetes?	5
Who gets gestational diabetes?	9
How do I find out if I have gestational diabetes?	10
What can I do if I have gestational diabetes?	11
What if I feel stressed?	12
What are the risks for baby?	13
What are the risks for me?	14
How do I keep healthy during my pregnancy?	15
I need to take insulin, what does this mean?	20
Hypoglycaemia (hypo)	21
What can I do to stay healthy after my baby is born?	22
Helpful contacts	24
Notes	26



- → Gestational diabetes is a type of diabetes that you can get when you are pregnant.
- Gestational diabetes is sometimes called GDM by health workers.
- → Gestational diabetes is when you have high blood glucose (sugar) levels during pregnancy.


When you eat or drink, your body breaks down carbohydrate foods into glucose, which then goes into your blood.




Your muscle cells need glucose to give your body energy to work, like how a car needs petrol to run.



Your pancreas makes insulin, which also goes into your blood. You need insulin to help the glucose move from your blood and into your muscles and cells.




Insulin acts like a key to unlock your muscles and cells and let the glucose in to be used for energy.




Your body knows when you have eaten and have glucose in your blood and it uses insulin to keep your blood glucose level balanced.

When you are pregnant your body makes hormones that help your baby grow. These normal pregnancy hormones block insulin from working properly, so your insulin keys can't open as many doors.



This means you need to make more insulin when you are pregnant.



Some women can't make enough insulin. If you can't make enough insulin the glucose builds up in your blood and you have gestational diabetes.





Who gets gestational diabetes?

→ All pregnant women can get gestational diabetes.



Women may be at greater risk of getting gestational diabetes if they:

- had gestational diabetes in a past pregnancy
- are over 30 years old
- have a family history of diabetes
- are an Aboriginal or Torres Strait Islander woman
- are an Asian woman or woman from other ethnic backgrounds
- are overweight.

How do I find out if I have gestational diabetes?



- → All pregnant women should have an Oral Glucose Tolerance Test to find out how their body responds to glucose.
- → The Oral Glucose Tolerance Test normally happens when your are 24–28 weeks pregnant, but your health worker might decide to do it sooner.

When you have the Oral Glucose Tolerance Test:

- You will be asked to not eat for eight hours before the test.
 This is called fasting.
- You will be asked to drink a very sweet drink and your blood will be tested several times.
- The test takes 2 hours, so take something to read while you are waiting.
- If your blood glucose is too high after the 2 hours, then you have gestational diabetes.

What can I do if I have gestational diabetes?



The best place to start is by visiting your health worker or clinic to get a health plan to help you look after yourself and your baby.

It is a good idea to take your partner, a family member or a friend with you to all your pregnancy appointments.

If you have gestational diabetes and do not follow the health plan from your health worker, you or your baby may become sick.



What if I feel stressed?

- → You may find managing your gestational diabetes hard. Many people feel stressed during this time.
- → Make sure you have lots of support from family and friends and talk to your health worker about how you feel.



What are the risks for baby?

It is important to manage your diabetes. If your blood glucose levels are too high during pregnancy then it may affect your baby's health.

It may lead to:

- baby growing bigger than he/she should. A large baby can make labour long and giving birth harder
- birth of your baby by caesarean
- baby being born too early

After the birth baby may have:

- trouble breathing and feeding
- low blood glucose levels
- jaundice (yellow skin and eyes)
- a higher risk of developing type 2 diabetes when they grow up

Baby will not be born with diabetes.



What are the risks for me?

If you have gestational diabetes:

- You are more at risk of high blood pressure during pregnancy.
- You are at greater risk of developing type 2 diabetes in the future.

Type 2 diabetes is very serious and can cause other long-term health problems.





It is important to look after yourself.


Make sure you go to all of your appointments. Take your partner, a family member or a friend to all of your appointments.

Managing your blood glucose levels during pregnancy will help keep you and your baby healthy.

Choosing healthy foods, doing exercise, taking medication and insulin and remembering to do your home blood glucose checking are all important.



Enjoy foods from each of the five food groups every day.



Have a yarn with your health worker about a healthy eating plan.

Drink lots of water



Don't drink alcohol



Stay away from sugary drinks.




Don't smoke



Alcohol and smoking can harm you and your baby. Talk to your health worker to get help to stop or reduce drinking and smoking.





Check your blood glucose levels regularly

- Follow your health worker's instructions on how often to check your blood glucose levels. Your doctor or diabetes educator will advise you on what blood glucose levels you should be aiming for.
- Talk to your health worker if you are worried about your blood glucose levels.
- If you feel unwell, talk to your health worker immediately as this can affect your blood glucose levels.

I need to take insulin, what does this mean?



Your doctor may recommend tablets or insulin injections to help you manage your gestational diabetes.

If you are taking insulin to manage your blood glucose levels, you could have hypoglycaemia (hypos). This is when your blood glucose level is low.

If you are taking insulin it is even more important to check your blood glucose levels carefully.

If you are taking insulin to avoid hypglycaemia (hypo):

- Never skip scheduled meals or snacks.
- Make sure you have hypo foods or treatments on you at all times, such as jelly beans.
- Check your blood glucose levels and check your insulin needs carefully.
- Keep your appointments and tell your health worker if you have a hypo.
- Tell people close to you the signs of a hypo and how to treat it.

Hypoglycaemia (hypo)

Hypoglycaemia (hypos) is when your blood glucose level drops below 4.0 mmol/L or when you begin to feel:

weak or tired	confused	headache
sweaty	shaky or dizzy	angry or moody
hungry	blurry vision	

If possible, check your blood glucose levels. If they are below 4.0 mmol/L then:

Hypo treatment

Step 1. Have:



5-7 jellybeans



1 glass of soft drink (not diet or sugar free)

OR



3 teaspoons of sugar dissolved in water

OR

OR

Step 2. Wait 10 minutes and check your blood glucose level. If it is below 4 mmol/L and you are not feeling better then go back to Step 1. If you are feeling better and your blood glucose is above 4 mmol/L have:



a sandwich



a glass of milk




a piece of fruit

What can I do to stay healthy after my baby is born?

Most of the time your blood glucose levels will go back to the healthy levels and your gestational diabetes will go away when the baby is born.

- You need to be checked six weeks after the baby is born to make sure you don't still have diabetes.
- You will need to see your clinic every one to two years to check if you have developed type 2 diabetes.



- Breastfeeding will help give your baby the best start. It can help stop
 baby becoming overweight and getting type 2 diabetes when they
 grow up. Breastfeeding can also help you to lose weight gained in
 pregnancy and may prevent you from getting type 2 diabetes later.
- Choose healthy foods, do regular exercise and maintain a healthy weight.
- If you are planning to become pregnant again make sure you talk to your health worker.



Helpful contacts

Diabetes WA Helpline

Our friendly diabetes educators are here to talk to you about diabetes and answer any questions you may have.



A diabetes educator is available 8:30 to 4:30 every week day.

Diabetes Telehealth for Country WA

Diabetes Telehealth provides diabetes education via videoconferencing technology to people in regional and rural WA.



Appointments are free and it is like having a face-to-face education session with a diabetes educator.

For more information or to book an appointment call 1300 001 880 or email telehealth@diabeteswa.com.au

Helpful contacts

National Diabetes Services Scheme

The National Diabetes Services Scheme (NDSS) delivers subsidised products, information and support services to people with diabetes across Australia.



For questions or to register please visit www.ndss.com.au/GD



Notes

Notes

My local clinic details:



Diabetes WA Aboriginal Health Team

Level 3, 322 Hay Street, Subiaco WA 6008

Open: 8.30am - 4.30pm weekdays

E: aboriginalhealth@diabeteswa.com.au

P: 1300 001 880 (cost of a local call)